

# Characteristics Of Oral Health In Early Onset Obesity

[ebook.ecog-obesity.eu/chapter-clinics-complications/characteristics-oral-health-early-onset-obesity](http://ebook.ecog-obesity.eu/chapter-clinics-complications/characteristics-oral-health-early-onset-obesity)



**Emmanuelle Noirrit-Esclassan**

**Mathieu Marty**

**Isabelle Bailleul-Forestier**

Dental university, Paediatric Department, Toulouse France

3 Chemin des maraîchers 31400 Toulouse, France

Corresponding author: I. Bailleul-Forestier

Email: [isabelle.bailleul-forestier@univ-tlse3.fr](mailto:isabelle.bailleul-forestier@univ-tlse3.fr)

The origins, and metabolic and physical outcomes of early onset of obesity are largely described in previous chapters. This section focuses on the oral cavity of children who are overweight or obese. Epidemiological and case-control studies have determined orofacial features specific to children of this population. Orofacial specificities in overweight or obese children are of various types. They could correspond to orofacial growth variations, increased risk of dental injuries, caries susceptibility, erosion, and periodontal defects.

## **Orofacial specificities**

Obese children have a normal to accelerated growth rate despite low growth hormone (GH) levels. Craniofacial growth and development are complex mechanisms regulated by interactions among genes, hormones, nutrients and epigenetic factors. They lead to the final craniofacial morphology. Patients with GH deficiency exhibit a smaller anterior and posterior cranial base, and small posterior facial and mandibular height [1]. In a Swedish craniofacial morphometric case-control study, obese adolescents showed increased mandibular length, prognathic jaws and a reduced upper anterior face height. Despite low GH levels, obese children have normal levels of insulin-like growth factor (IGF-1). Craniofacial growth may be more dependent on free circulating IGF-1 [2]. So obese subjects exhibit an increase of some craniofacial parameters.

In Iranian and US populations, children who were overweight or obese, after being adjusted for age and gender, showed accelerated dental development whereas they did not have accelerated skeletal maturation [3,4]. According to the BMI classification, Costacurta et al. observed no statistical differences for chronological, dental and skeletal age between normal weight and obese children. However when the Fat Mass % was taken into account (McCarthy classification) the skeletal-dental age always increased with respect to the chronological age [5]. For orthodontic treatment in overweight or obese children, determination of skeletal maturation and dental development is important and the specificities of the jaw growth should be considered. Additionally, an earlier onset of puberty has been reported in children with higher percentile [6]. The onset of puberty conditions the beginning of orthodontic treatment. So these findings may be of interest in orthodontic procedures. Young obese subjects may need a different orthodontic treatment plan in comparison to normal weight patients of the same age. However, no protocol currently exists for the orthodontic treatment process of overweight or obese children.

In children, obstructive sleep apnea syndrome (OSAS) belongs to a broad spectrum of respiratory disorders that occur during sleep, and includes primary snoring and upper airway resistance syndrome. OSAS occurs more frequently and may be more severe in children and adolescents who are overweight or obese compared with lean children. [7]. In obese adolescents, the enlargement of the upper airway soft tissue structures, the increased size of the pharyngeal lymphoid tissue, and dysfunctional breathing contribute to OSAS. These children exhibit mouth breathing, narrower maxilla, open bite and lateral posterior crossbite. Adenotonsillectomy should be considered as the first treatment option for OSAS in obese adolescents. But it does cure half of them [7]. However, obesity at the time of diagnosis is a major risk for persisting OSAS after treatment, regardless of the severity of initial disease [8]. Emerging dental treatment options, such as rapid maxillary expansion, oral appliances and osteogenesis distraction may soon serve as viable or adjunctive treatment alternatives to adenotonsillectomy. These treatments help to normalize the craniofacial and dentofacial morphology, and to modify tongue posture and the airway, thus leading to restoration of a normal trajectory of facial growth and development in children [9].

## Dental injuries

An increase in body fatty mass reduces postural stability by moving the centre of gravity. Moreover, obese patients have a functional limitation of the articulations [10]. An increased prevalence of bone fractures is detected among obese patients, a tendency that could be linked with dental traumatism [11,12].

Some cross-sectional studies have tried to clarify this relation, but various biases can influence the results (cf Table 1). The limits establishing obesity fluctuate, and obesity was measured at the time of the study but not at the time of the trauma [18]. The evaluation of dental traumatism was only visual (without any radiography or vitality test), so this risk was undervalued [16,18]. The variability of these features could explain the heterogeneity of the results. Correa-Faria and Petti (2015) carried out a meta-analysis starting from 17 studies, 13 of them concerning young adolescents (12 years old on average) [25]. They concluded that a link existed between overweight/obesity and dental traumatism, although the force of this association was less strong than for other risk factors, such as the incisor overjet. The risk of traumatism was increased by 22% if the patient was obese, whereas the risk was 131% in the presence of an incisive overjet [25]. As dental traumatism has increased prevalence among obese patients (28.6% of traumatism compared with 25.5% in slim children, Correa-Faria and Petti's (2015) meta-analysis suggests a causal association (and not only a statistical association) between obesity and dental traumatism [24].

Prevention of dental traumatism involves orthodontic interception of malocclusion and the subject wearing a mouth guard when practising sports. An additional reason for managing excess weight is to avoid such dental injuries.

## Caries risk

Both obesity and dental caries are important health issues with multifactorial aspects. They have become increasingly prevalent challenges to public health. Factors affecting the onset of carious lesions include oral hygiene, diet composition and frequency, salivary immunoglobulins, bacterial load, fluoride intake and socioeconomic status [26]. Overweight, obesity and dental caries are concomitant in many population groups, probably because of the common confounding risk factors such as socioeconomic status, high sugar intake frequency, cariogenic diet, and sedentary life style. Even the notion that they have a cause–effect relationship should be viewed with caution.

A number of studies have linked consumption of sweetened beverages with increased energy intake, obesity [27] and dental decay [28]. Some studies in primary, mixed and permanent dentitions report an association between obesity and dental caries [29, 30]. However the data collected in cross-sectional US surveys (NHANES, Third National Health and Nutrition Examination Survey (NHANES III)) showed no evidence that overweight children were at increased risk for dental caries [31,32]. In 2006, Kantowics conducted a first systematic review (1984 to 2004) reporting that only one of seven cross-sectional studies with children showed an association between obesity and dental caries [33]. A newer systematic review and meta-analysis quantified the relationship between obesity and dental caries in children using a systematic approach and considering the literature over a longer period (1980 to 2010) [34]. Overall, a significant relationship between childhood obesity and dental caries (effect size = 0.104,  $P = 0.049$ ) was found in the studies using standardized measures for assessing child

obesity such as BMI-for-age centiles (effect size = 0.189, 95% CI: 0.060–0.318,  $P = 0.004$ ) or IOTF cut-offs (effect size = 0.104, 95% CI: 0.060–0.180,  $P = 0.008$ ). These studies, using standard assessment of child obesity analyses by dentition type, showed a significant association between obesity and dental caries in permanent (effect size = 0.198, 95% CI: 0.046–0.350,  $P = 0.011$ ) but not in primary dentitions. Evaluations between normal-weight and obese children in industrialized countries (IC) showed a significant relationship between obesity and caries (effect size = 0.122, CI = 0.047–0.197,  $P = 0.001$ ) in contrast to those from non-industrialized countries (NIC) (effect size = 0.079, CI = !0.106 to 0.264,  $P = 0.264$ ) [34]. Mixed results for the relationship between obesity and dental caries could be explained by the use of standardized or non-standardized BMI, different growth rates among children of similar age [35], or genetic susceptibility to decay [36].

Based on the concept that diet, and in particular sugar, contributes to the development of dental caries and overweight, dental personnel have been suggested to be one of the cornerstones in weight counselling [37].

Another increasing dental trouble, dental erosion, could be statistically observed in young subjects above the 98<sup>th</sup> percentile for BMI. Erosion was more severe with a larger area affected than in normal weight children [38]. Children with gastroesophageal reflux disease (GERD) are more likely to have erosion defects. Moreover in children and adolescent, obesity is related to increased odds of GERD [39]. So dental erosion should be carefully sought in overweight or obese children to set up preventive and aetiological treatment (fluoride).

## **Periodontal diseases in children and adolescents.**

Gingivitis and periodontitis are the two major forms of periodontal diseases, which are defined as inflammatory diseases affecting the periodontium. According to the American Academy of Pediatric Dentistry, gingivitis is an inflammation of the gingiva that does not result in clinical attachment loss. Periodontitis is characterized by an inflammation of the gingiva and by loss of connective tissue attachment and alveolar bone [40]. Untreated, it may lead to tooth loss. The primary aetiology of both diseases is bacterial dental biofilm that adheres to the tooth surface. Clinically, it is characterized by gum inflammation with oedema, bright red or purple colour and bleeding on probing [41]. The prevalence of periodontal diseases in children is lower than in adult patients [42]. However, gingivitis in childhood is considered as a risk factor for attachment loss in adult life [43].

Statistical link between obesity and periodontal diseases among children and adolescents

An exhaustive review of this link in current databases found 10 clinical trials and meta-analysis fitting the aims. The results are summarized in Table 2. Most of the authors observed that obesity was linked to periodontal diseases. First of all, a higher Plaque Index was observed in obese adolescents, who had poorer toothbrushing habits than individuals of normal weight [45,50]. Clinically, a simple gum inflammation (gingivitis) was frequently found, but attachment loss was not [44; 45; 49; 50]. However, Cinar et al. and Modéer et al. found greater pocket depth in obese patients [52;53]. Only one study did not show a relationship between obesity and periodontal diseases but the clinical examination was unclear [47].

Despite the great heterogeneity in the clinical indicators used, and in the groups of patients, a statistical association could be found between obesity and clinical gum inflammation. Biological markers of periodontal diseases were also more commonly found in patients suffering from metabolic syndrome or obesity. However, no evidence could be found to link obesity and bone loss. Moreover,

as with normal-weight individuals, the periodontal risk factors increased with age, and the adolescents and young adults were more affected than young children [44, 45].

The exact effect of obesity on the periodontium is not clearly understood. The most common inflammation markers used to validate the presence of periodontitis are interleukin-1" (IL-"), interleukin-8 (IL-8) and tumour necrosis factor # (TNF-#). TNF# plays a significant role in the pathogenesis and development of periodontal diseases and is found in the crevicular fluid of patients suffering from periodontitis [55]. IL-8 is a neutrophil chemoattractant produced during the inflammatory process [56]. These factors were commonly found to be linked with obesity in the paediatric samples of two studies [48; 53]. Obesity contributes to a pro-inflammatory environment by producing pro-inflammatory cytokines. For Katz et al., such cytokines, associated with other local factors, may contribute to gum disease in children and adolescents [51]. Moreover, it has been proved that some chronic infections may contribute to obesity development [57]. Although a recent review demonstrates that factors such as lifestyle and type 2 diabetes may confound the linkage between periodontal disease and obesity [58], physicians facing an obese child or adolescent may refer him to a paediatric dentist for a specialized consultation.

Clinical implications for oral prevention and treatment of child or adolescent with early onset of obesity

For the dental practitioner, the clinical examination of an overweight child or adolescent should evaluate oral functions in order to detect possible mouth breathing, and cross or open bite leading to mastication deficiencies. Any orthodontic treatment plan should consider variations in facial growth and dental maturation in children or adolescents with early onset obesity.

This examination routinely includes a caries risk assessment and a community periodontal index [59]. In case of doubt, a complete examination, including periodontal charting and radiological status should be performed. Conversely, the medical follow up by a physician of a subject with early onset overweight or obesity should include a dental examination.

## Tables

**Table 1:** Review of link between obesity and dental injuries among children and adolescents

Author (year)	Country	Population: N	Age	BMI	Obesity and dental trauma association
Petti (1997) [13]	Italy	938	6-11	> 97th percentile	Obesity significantly increased the risk of trauma: 1/3 for obese and 1/5 for non-obese children.
Nicolau (2001) [14]	Brazil	652	13	BMI > 85th percentile	Overweight children had 1.93 greater risk of dental trauma
Soriano (2009) [15]	Brazil	1046	12	> 97th percentile (NCHS reference table)	Obese patients underwent 1.84 times more trauma than non-obese but “obesity” factor was not linked with trauma risk
Artun (2009) [16]	Kuwait	1583	13-14	>30	No difference of risk between 3 BMI groups: >18.5, <18,5 and <30, >30
Damé-Teixeira (2013) [17]	Brazil	1528	12	BMI > 85th percentile	No significant relationship between obesity and trauma
Al-Bajjali (2014) [18]	Jordan	1015	12	BMI>24.2 boys, BMI>25.2 girls (Centre for Disease Control and Prevention)	No significant relationship between obesity and trauma
Basha (2015) [19]	India	764	13	Overweight: 20.3-24.9 boys, 22.7-26.8 girls Obese: >25 boys, >26.9 girls (criteria of Indian Association of Pediatrics)	Risk of suffering trauma 2.78 time higher in obese and overweight children
Tapias (2003) [20]	Spain	388	10	BMI > 85th percentile	No relation between overweight and dental traumatism
Vijaykumar (2013) [21]	India	858	10-12 (10.81+/-0.43)	BMI > 95th percentile (Centre for Disease Control and Prevention)	Significant association between trauma and overweight: 3.85 greater risk of trauma in overweight patients, of either sex
Goettems (2014) [22]	Brazil	1203	8-12	BMI > 85th percentile	1.2 more risk of dental trauma in obese and overweight subjects
Martins (2012) [23]	Brazil	590	7-14	BMI > 85th percentile	No statistical link in multi variable analysis

**Table 2:** Review of link between obesity and periodontal diseases among children and adolescents

Author, (year)	Study type	Patients / control	Obesity and periodontal disease association
Galkina IuV et al. 2015 [44]	Observational study	168 obese adolescents / no control	Yes
Markovic D et al. 2014 [45]	Cross-sectional study	442 patients aged 6-18 years	Yes, association with Gingival Index
Scorzetti L et al. 2013 [46]	Case control study	44 obese patients aged 7-13 years / 59 controls aged 8-10 years	Yes concerning the risk factors but not attachment loss
Nascimento GG et al. 2013 [47]	Cross-sectional study	419 obese patients aged 8-12 years / 792 controls aged 8-12 years	No correlation between obesity and gingivitis but obese boys have higher risk of gingivitis.
Kâ K et al. 2013 [48]	Cross-sectional study	175 obese patients, 25 of whom had metabolic syndrome aged 8-10 / 273 patients aged 8-10	Patients with metabolic syndrome show higher inflammation factor and gum bleeding, especially in boys.
Fadel HT et al. 2014 [49]	Case-control study	27 obese adolescents / 28 controls	Yes, more gingivitis but no increase in depth of periodontal pockets.
Franchini R et al. 2011 [50]	Case-control study	47 obese and 48 controls aged 10-17 years	Yes higher gingival index among obese patients.
Mod�er T et al. 2011 [52]	Case-control study	52 obese adolescents/ 52 controls	Yes, more gum inflammation, more periodontal pockets, greater periodontal pocket depth, but no bone loss.
Cinar AB et al. 2011 [53]	Cross-sectional study	611 patients	Yes, more gum inflammation and greater pocket depth.
Reefes AF et al 2006 [54]	Case control study	111 adolescents aged 13-21 years with at least one periodontal pocket. 2341 controls.	An increased risk of periodontal disease is related to waist circumference and weight.

## References

- 1 Pirinen S, Majurin A, Lenko HL, Koski K. Craniofacial features in patients with deficient and excessive growth hormone. *J Craniofac Genet Dev Biol.* 1994;14:144-52.
- 2 Ohrn K, Al-Kahlili B, Huggare J, Forsberg CM, Marcus C, Dahllöf G. Craniofacial morphology in obese adolescents. *Acta Odontol Scand.* 2002;60:193-7.
- 3 Hilgers KK, Akridge M, Scheetz JP, Kinane DE. Childhood obesity and dental development. *Pediatr Dent.* 2006;28:18-22.
4. Hedayati Z, Khalafinejad F. Relationship between Body Mass Index, Skeletal Maturation and Dental Development in 6- to 15- Year Old Orthodontic Patients in a Sample of Iranian Population. *J Dent (Shiraz).* 2014;15:180-6.
5. Costacurta M, Sicuro L, Di Renzo L, Condò R, De Lorenzo A, Docimo R. Childhood obesity and skeletal-dental maturity. *Eur J Paediatr Dent.* 2012;13:128-32.
6. Sandhu J, Ben-Shlomo Y, Cole TJ, Holly J, Davey Smith G. The impact of childhood body mass index on timing of puberty, adult stature and obesity: a follow-up study based on adolescent anthropometry recorded at Christ's Hospital (1936-1964). *Int J Obes.* 2006;30(1):14-22.
7. Mathew JL, Narang I. Sleeping too close together: obesity and obstructive sleep apnea in childhood and adolescence. *Paediatr Respir* 2014; 15: 211-8.
8. O'Brien LM, Sitha S, Baur LA, Waters KA. Obesity increases the risk for persisting obstructive sleep apnea after treatment in children. *Int J Pediatr Otorhinolaryngol.* 2006;70:1555-60.
9. Ngiam J, Cistulli PA. Dental treatment for paediatric obstructive sleep apnea. *Paediatr Respir Rev.* 2015;16:174-81.
10. Maffiuletti NA, Agosti F, Proietti M, Riva D, Resnik M, Lafortuna CL, Sartorio A. Postural instability of extremely obese individuals improves after a body weight reduction program entailing specific balance training. *J Endocrinol Invest.* 2005;28:2-7.
11. Wills M. Orthopedic complications of childhood obesity. *Pediatr Phys Ther.* 2004;16:230-5.
- 12 Dimitri P, Bishop N, Walsh JS, Eastell R. Obesity is a risk factor for fracture in children but is protective against fracture in adults: a paradox. *Bone.* 2012;50:457-66.
- 13 Petti S, Cairella G, Tarsitani G. Childhood obesity: a risk factor for traumatic injuries to anterior teeth. *Endod Dent Traumatol.* 1997;13:285-8
- 14 Nicolau B, Marcenes W, Sheiham A. *Dent Traumatol.* 2001;17:213-7. Prevalence, causes and correlates of traumatic dental injuries among 13-year-olds in Brazil.
15. Soriano EP, Caldas Ade F Jr, De Carvalho MV, Caldas KU. Relationship between traumatic dental injuries and obesity in Brazilian schoolchildren. *Dent Traumatol.* 2009;25:506-9.



16. Artun J, Al-Azemi R. Social and behavioral risk factors for maxillary incisor trauma in an adolescent Arab population. *Dent Traumatol*. 2009;25:589-93.
17. Damé-Teixeira N, Alves LS, Susin C, Maltz M. Traumatic dental injury among 12-year-old South Brazilian schoolchildren: prevalence, severity, and risk indicators. *Dent Traumatol*. 2013;29:52-8.
18. Al-Bajjali TT, Rajab LD. Traumatic dental injuries among 12-year-old Jordanian schoolchildren: an investigation on obesity and other risk factors. *BMC Oral Health*. 2014;14:101.
19. Basha S, Mohammad RN, Swamy HS. Incidence of dental trauma among obese adolescents--a 3-year-prospective study. *Dent Traumatol*. 2015;31:125-9.
20. Tapias MA, Jiménez-García R, Lamas F, Gil AA. Prevalence of traumatic crown fractures to permanent incisors in a childhood population: Móstoles, Spain. *Dent Traumatol*. 2003;19:119-22.
21. Vijaykumar S, Guna Shekhar M, Vijayakumar R. Traumatic Dental Injuries and Its Relation to Overweight among Indian School Children Living in an Urban Area. *J Clin Diagn Res*. 2013;7:2631-3.
22. Goettems ML, Torriani DD, Hallal PC, Correa MB, Demarco FF. Dental trauma: prevalence and risk factors in schoolchildren. *Community Dent Oral Epidemiol*. 2014;42:581-90.
23. Martins VM, Sousa RV, Rocha ES, Leite RB, Paiva SM, Granville-Garcia AF. Dental trauma among Brazilian schoolchildren: prevalence, treatment and associated factors. *Eur Arch Paediatr Dent*. 2012;13:232-7.
24. Corrêa-Faria P, Petti S. Are overweight/obese children at risk of traumatic dental injuries? A meta-analysis of observational studies. *Dent Traumatol*. 2015;31:274-82.
25. Petti S. Over two hundred million injuries to anterior teeth attributable to large overjet: a meta-analysis. *Dent Traumatol*. 2015;3:1-8.
26. Caufield PW, Li Y, Dasanayake A. Dental caries: an infectious and transmissible disease. *Compend Contin Educ Dent*. 2005; 26:10-6.
27. Hu FB. Resolved: there is sufficient scientific evidence that decreasing sugar-sweetened beverage consumption will reduce the prevalence of obesity and obesity-related diseases. *Obes Rev*. 2013;14:606-19.
28. Marshall TA, Levy SM, Broffitt B, Warren JJ, Eichenberger-Gilmore JM, Burns TL et al. Dental caries and beverage consumption in young children. *Pediatrics* 2003;112:e184-91.
29. Alm A, Fahraeus C, Wendt LK, Koch G, Andersson-Gare B, Birkhed D. Body adiposity status in teenagers and snacking habits in early childhood in relation to approximal caries at 15 years of age. *International Journal of Paediatric Dentistry* 2008;18:189-196.
30. Gerdin EW, Angbratt M, Aronsson K, Eriksson E, Johansson I. Dental caries and body mass index by socio-economic status in Swedish children. *Community Dent Oral Epidemiol* 2008;36:459-65.

31. Kopycka-Kedzierawski DT, Auinger P, Billings RJ, Weitzman M. Caries status and overweight in 2- to 18-year-old US children: findings from national surveys. *Community Dent Oral Epidemiol* 2008;36:157–67.
32. Macek MD, Mitola DJ. Exploring the association between overweight and dental caries among US children. *Pediatr Dent* 2006;28:375–80.
33. Kantovitz KR, Pascon FM, Rontani RM, Gaviao MB. Obesity and dental caries—a systematic review. *Oral Health Prev Dent* 2006;4:137–44.
34. Hayden C, Bowler JO, Chambers S, Freeman R, Humphris G, Richards D, Cecil JE. Obesity and dental caries in children: a systematic review and meta-analysis. *Community Dent Oral Epidemiol*. 2013;41:289-308.
35. Cole TJ, Bellizzi MC, Flegal KM, Dietz WH. Establishing a standard definition for child overweight and obesity worldwide: international survey. *BMJ*. 2000;320:1240-3.
36. Opal S, Garg S, Jain J, Walia I. Genetic factors affecting dental caries risk. *Aust Dent J*. 2015;60:2-11.
37. Lempert SM, Froberg K, Christensen LB, Kristensen PL, Heitmann BL. Association between body mass index and caries among children and adolescents. *Community Dent Oral Epidemiol*. 2014;42:53-60.
38. Tong HJ, Rudolf MC, Muyombwe T, Duggal MS, Balmer R. An investigation into the dental health of children with obesity: an analysis of dental erosion and caries status. *Eur Arch Paediatr Dent*. 2014;15:203-10
39. Koebnick C, Getahun D, Smith N, Porter AH, Der-Sarkissian JK, Jacobsen SJ. Extreme childhood obesity is associated with increased risk for gastroesophageal reflux disease in a large population-based study. *Int J Pediatr Obes*. 2011; 6:e257-63.
40. American Academy of Pediatric Dentistry. Treatment of Plaque-induced Gingivitis, Chronic Periodontitis, and Other Clinical Conditions. *Clinical Guideline*, 2014;36, 360–369.
41. American Academy of Periodontology. The pathogenesis of periodontal diseases (position paper). *J Periodontol* 1999;70:457-70.
42. American Academy of Periodontology. Diagnosis of Periodontal Diseases (position paper). Chicago, Ill: The American Academy of Periodontology; 1995.
43. Albandar JM, Tinoco EMB. Global epidemiology of periodontal diseases in children and young persons. *Periodontol* 2000. 2002;29:153-176.
44. Oh Tj, Eber R, Wang HL. Periodontal diseases in the child and adolescent. *J Clin Periodontol* 2002;29: 400-410.

45. Galkina IuV, Gavrilova OA, Piekalnits Iia, Dianov OA. Dental status in children and adolescents diagnosed with obesity. *Stomatologia (Mosk)*. 2015;94:57-8.
46. Markovic D, Ristic-Medic D, Vucic V, Mitrovic G, Nikolic Ivosevic J, Peric T, Karadzic I. Association between being overweight and oral health in Serbian schoolchildren. *Int J Paediatr Dent*. 2015; 25:409-17.
47. Scorzetti L, Marcattili D, Pasini M, Mattei A, Marchetti E, Marzo G. Association between obesity and periodontal disease in children. *Eur J Paediatr Dent*. 2013;14:181-4.
48. Nascimento GG, Seerig LM, Vargas-Ferreira F, Correa FO, Leite FR, Demarco FF. Are obesity and overweight associated with gingivitis occurrence in Brazilian schoolchildren? *J Clin Periodontol*. 2013;40(12):1072-8.
49. Kâ K, Rousseau MC, Lambert M, Tremblay A, Tran SD, Henderson M, Nicolau B. Metabolic syndrome and gingival inflammation in Caucasian children with a family history of obesity. *J Clin Periodontol*. 2013;40:986-93.
50. Fadel HT, Pliaki A, Gronowitz E, Mårild S, Ramberg P, Dahlèn G, Yucel-Lindberg T, Heijl L, Birkhed D. Clinical and biological indicators of dental caries and periodontal disease in adolescents with or without obesity. *Clin Oral Investig*. 2014;18:359-68.
51. Franchini R, Petri A, Migliario M, Rimondini L. Poor oral hygiene and gingivitis are associated with obesity and overweight status in paediatric subjects. *J Clin Periodontol*. 2011;38:1021-8.
52. Katz J, Bimstein E. Periodontal disease and obesity in children. *Isr Med Assoc J*. 2010;12:775-6.
53. Modéer T, Blomberg C, Wondimu B, Lindberg TY, Marcus C. Association between obesity and periodontal risk indicators in adolescents. *Int J Pediatr Obes*. 2011;6(2-2):e264-70.
54. Cinar AB, Murtomaa H. Interrelation between obesity, oral health and life-style factors among Turkish school children. *Clin Oral Investig*. 2011;15:177-84.
55. Reeves AF, Rees JM, Schiff M, Hujuel P. Total body weight and waist circumference associated with chronic periodontitis among adolescents in the United States. *Arch Pediatr Adolesc Med*. 2006;160:894-9.
56. Grauballe MB, Østergaard JA, Schou S, Flyvbjerg A, Holmstrup P. Effects of TNF- $\alpha$  blocking on experimental periodontitis and type 2 diabetes in obese diabetic Zucker rats. *J Clin Periodontol*. 2015; 42:807-16.
57. Lappin DF, Robertson D, Hodge P, et al. Evaluation of Serum Glycated Haemoglobin, IL-6, IL-8 and CXCL5 in T1DM With and Without Periodontitis and Effects of Advanced Glycation End Products and Porphyromonas Gingivalis Lipopolysaccharide on IL-6, IL-8 and CXCL5 Expression by Oral Epithelial Cells and Monocytes. *J Periodontol*. 2015; 7:1-20.
58. Dhurandhar NV, Bailey D, Thomas D. Interaction of obesity and infections. *Obes Rev*. 2015;16: 1017-19.

59. Olsen I. From the Acta Prize Lecture 2014: the periodontal-systemic connection seen from a microbiological standpoint. *Acta Odontol Scand.* 2015;73:563-8.
60. Clerehugh V, Tugnait A. Diagnosis and management of periodontal diseases in children and adolescents. *Periodontol 2000.* 2001;26:146-168.

## ~ About the Authors ~



**Pr Isabelle Bailleul-Forestier, DDS, PhD**, is a professor and head of the department of Pediatric Dentistry at Dental Faculty, Toulouse University, France.

The topic of her PhD was the expression of orodental signs of systemic and rare diseases. She is Visiting Professor, at the Centre of Human Genetics, Leuven, Belgium She contributes to clarify the relationship between obesity and dental caries in sever obese adolescents.



**Dr Emmanuelle Noirrit-Esclassan, DDS, PhD** is senior lecturer in the department of pediatric dentistry at Children's hospital in Toulouse, France. In clinics, she is taking care of children with severe early-childhood caries and special care children under sedation.



**Dr Mathieu Marty, DDS, Msc**, is an assistant professor in the department of paediatric dentistry at Toulouse University Hospital. He is involved in the paediatric oral mucosa pathology consultation.

His main research topic concerns pediatric periodontitis. Particularly, his work intends to develop diagnosis tools and periodontal treatment procedure for patient in medically underserved area with Pr J Palmieri (Blacksburg, USA).

### ~ How To Use This article ~

You are **free to use, share and copy this content** by quoting this article as follow:

*Noirrit-Esclassan E, Marty M, Bailleul-Forestier I (2015). XXX. In M.L. Frelut (Ed.), The ECOG's eBook on Child and Adolescent Obesity. Retrieved from [ebook.ecog-obesity.eu](http://ebook.ecog-obesity.eu)*

Also make sure to **give appropriate credit** when using this content. Please visit [ebook.ecog-obesity.eu/terms-use/summary/](http://ebook.ecog-obesity.eu/terms-use/summary/) for more information.

### ~ Final Word ~

Thank you for reading this article.

If you have found this article valuable, please share it with someone that will be interested in.

Also make sure to visit [ebook.ecog-obesity.eu](http://ebook.ecog-obesity.eu) to read and download more childhood obesity-related articles.